

ranging from  $9170 \pm 70$  BP (10,350-9,980 Cal BP age – AA18516) for a skull from Newburgh, Fife to  $3315 \pm 55$  bp (3690-3390 cal BP – AA-18517) for skull from Galloway. The latter is one of the most recent dates for Britain, suggesting that it survived until at least the early Bronze Age in Scotland (Yalden & Kitchener, 2008). A similar date was recorded for a skeleton from Charterhouse Warren Farm, Somerset (Burleigh & Clutton-Brock, 1977). However, the Ardyge Farm specimen is the oldest recorded post-glacial date for an aurochs in Scotland, and demonstrates that this species was an early post-glacial coloniser.



a.



b.

**Fig. 1.** Dorsal (a.) and ventral (b.) views of the horn cores of an aurochs, *Bos primigenius*, from Ardyge Farm, Morayshire (Neil McLean, National Museums Scotland).

The horn cores are available for viewing by appointment at Moray Estates (013096 72213 or admin@medeo.co.uk).

## REFERENCES

- Burleigh, R. & Clutton-Brock, J. (1977). A radiocarbon date for *Bos primigenius* from Charterhouse Warren Farm, Mendip. *Proceedings of the University of Bristol Speleological Society* 14(3), 225-257.
- Kitchener, A.C. & Bonsall, C. (1999). Further AMS radiocarbon dates for extinct Scottish mammals. *Quaternary Newsletter* No. 88, 1-10.

- Lowc, J.J. and Walker, M.J.C. (1997). *Reconstructing Quaternary environments*. 2nd edition Prentice Hall, Harlow.
- Yalden, D. (1999). *The history of the British mammals*. Poyser, London.
- Yalden, D.W. & Kitchener, A.C. (2008). History of the fauna. Pp. 17-31. In: Harris, S. and Yalden, D.W. (editors). *Mammals of the British Isles: Handbook, 4<sup>th</sup> edition*. The Mammal Society, Southampton.

## The rare green alga *Pediastrum privum* (Chlorophyta, Sphaeropleales) in a Scottish kettle loch: new to British freshwaters

<sup>1</sup>Pauline Lang, Jan Krokowski, Nicole Ross & Ross Doughty

Scottish Environment Protection Agency, 5 Redwood Crescent, Peel Park, East Kilbride, G74 5PP, Scotland, UK

<sup>1</sup>E-mail: pauline.lang@sepa.org.uk

*Pediastrum* is a widely-distributed genus of green alga characteristically consisting of disc-shaped colonies or 'coenobia', assembled from at least four inter-connecting cells (Komárek & Jankovská, 2001).

Many species belonging to the genus are common constituents of lake phytoplankton communities, though *Pediastrum privum* (Printz) Hegewald [= *Stauridium privum* (Printz) Hegewald in Buchheim *et al.*, 2005] is notably rare (Komárek & Jankovská, 2001; Tsarenko & John, 2011). There appear to be no published records from Britain. Sporadic lake phytoplankton and sub-fossil sediment finds from Europe, reflect a sparse scattering of *Pediastrum privum*, confined mostly to temperate and sub-arctic latitudes of the northern hemisphere (Hegewald & Schnepf, 1979; Komárek & Jankovská, 2001; Geriš, 2004; Kowalska & Wołowksi, 2010). By comparison, the close phylogenetic relative *Pediastrum tetras* (Ehrenberg) Ralfs [= *Stauridium tetras* (Ehrenberg) Hegewald in Buchheim *et al.*, 2005] displays a cosmopolitan distribution (Komárek & Jankovská, 2001).

Freshwater phytoplankton communities are important indicators of the biointegrity of standing waters and are therefore used by the Scottish Environment Protection Agency (SEPA) to assess the ecological status of around 80 freshwater lochs in Scotland. Phytoplankton samples are collected at varying frequencies, but at a minimum are taken three times a year between July and September. Sub-samples of phytoplankton (preserved in Lugol's iodine) are examined using an inverted microscope and analysed according to standard

procedures with counts of approximately 400 individuals (Brierley *et al.*, 2007; CEN, 2004 & 2008).

Low abundances (typically 1-5 coenobia, comprising both four- and eight-cells, per 100 ml sub-sample) of *Pediastrum privum* were found in phytoplankton samples collected from Loch Kinord during 2009–2011. Loch Kinord is a small kettle lake located in Aberdeenshire, Scotland (NGR: NO 44150 99388). The loch, formed by glacial retreat approximately 10,000 years ago, has an area of c. 0.8 km<sup>2</sup>, is shallow (mean depth <2 m) and is characterized by relatively low alkalinity (annual mean 10.7 mg L<sup>-1</sup> as CaCO<sub>3</sub> over 2009-11) and mesotrophic water chemistry (annual mean total phosphorus (TP) concentration 19.9 µg L<sup>-1</sup> over 2009-11). A palaeolimnological study using fossil diatoms implied that eutrophication has driven water quality in Loch Kinord slightly away from its reference state (Bennion *et al.*, 2004).

*Pediastrum privum* has appeared consistently in the phytoplankton community of Loch Kinord since 2009. This is the first known documented record of *Pediastrum privum* in British freshwaters. Previously, this uncommon species may have gone unnoticed or been misidentified due to its inconspicuous size and general unfamiliarity to UK taxonomists. The coenobia of *Pediastrum privum* (Figs 1a-d) morphologically resemble *Pediastrum tetras* (Figs 2a-d), in terms of their relatively small diameter (usually 15–25 µm). However, it is possible to separate the two species by comparison of the outer cell wall structure, which is weakly concave (central depression) in *P. privum* and distinctly notched (central incision) in *P. tetras* (Komárek & Jankovská, 2001; Kowalska & Wołowksi, 2010; Tsarenko & John, 2011).

*Pediastrum privum* has been recorded mostly from European waterbodies including Norway (Printz, 1914), Finland (Hegewald & Schnepf, 1979), Poland (Pelechaty *et al.*, 2007; Kowalska & Wołowksi, 2010), Russia (Jankovská & Komárek, 2000), Slovakia


(Hindák & Hindáková, 2008), and the Czech Republic (Gerš, 2004), though the WISER phytoplankton database ([www.wiser.eu](http://www.wiser.eu)) may also contain previously undocumented localities. Other reports exist from the USA (Smith, 1920; Prescott, 1962), as well as more recently from Korea (An *et al.*, 1999), Spain (Negro *et al.*, 2000) and Canada (Hindák & Hindáková, 2008). Collectively, observations suggest that *Pediastrum privum* occurs discretely in oligotrophic and/or dystrophic freshwaters (Jankovská & Komárek, 2000; Komárek & Jankovská, 2001). However, some accounts suggest it is also capable of occupying nutrient-enriched habitats (An *et al.*, 1999), typically associated with *P. tetras* (Komárek & Jankovská, 2001), which makes its restricted distribution difficult to explain (Kowalska & Wołowksi, 2010). Morphological plasticity (variation between the 4- and 8-celled life cycle stages) has been related to environmental nutrient concentrations or zooplankton predation in *Pediastrum tetras* (Rojo *et al.*, 2008), and though fully described (Hegewald & Jeon, 2000) is as yet inadequately understood for *P. privum*. More research is required to establish the ecological requirements of *Pediastrum privum* and the reasons for its apparent rarity.

#### ACKNOWLEDGEMENTS

Thanks especially to Prof David John (Natural History Museum London) and Prof Brian Whitton (University of Durham) for formally verifying the identity of *Pediastrum privum*. We are grateful to Dr Elizabeth Haworth (Freshwater Biological Association) for confirming that no UK records of *P. privum* pre-existed in the Fritsch Collection (documented only from Norway, Europe and North America). We are also appreciative to Dr Kevin Murphy (University of Glasgow) for commenting on an earlier version of the manuscript.



**Fig. 1a.** Photo-micrograph of *Pediastrum privum* 4-celled coenobium (x630 magnification) preserved in Lugol's Iodine.




**Fig. 1b.** Illustration of *Pediastrum privum* 4-celled coenobium.





**Fig. 1c.** Photo-micrograph of *Pediastrum privum* 8-celled coenobium (x630 magnification) preserved in Lugol's Iodine.




10  $\mu$ m

**Fig. 1d.** Illustration of *Pediastrum privum* 8-celled coenobium.




**Fig. 2a.** Photo-micrograph of *Pediastrum tetras* 4-celled coenobium (x630 magnification) preserved in Lugol's Iodine.




10  $\mu$ m

**Fig. 2b.** Illustration of *Pediastrum tetras* 4-celled coenobium.



**Fig. 2c.** Photo-micrograph of *Pediastrum tetras* 8-celled coenobium (x630 magnification) preserved in Lugol's Iodine.



10  $\mu$ m

**Fig. 2d.** Illustration of *Pediastrum tetras* 8-celled coenobium.

## REFERENCES

- An, S.S., Hegewald, E. & Jeon, S.L. (1999) *Pediastrum privum* (Printz) Hegewald new to Korea, *Algae*, 14, 83 – 85.
- Bennion, H., Fluin, J. & Simpson, G.L. (2004) Assessing eutrophication and reference conditions for Scottish freshwater lochs using subfossil diatoms, *Journal of Applied Ecology*, 41, 124 – 138.
- Brierley, B., Carvalho, L., Davies, S. & Krokowski, J. (2007) *Guidance on the quantitative analysis of phytoplankton in freshwater samples*, 24 pp. [In Carvalho, L., Dudley, B., Dodkins, I., Clarke, R., Jones, I., Thackeray, S., and Maberly, S. (2007) *Phytoplankton Classification Tool (Phase 2)*, Final Report, Project WFD80, SNIFFER, Edinburgh].
- Buchheim, M., Buchheim, J., Carlson, T., Braband, A., Hepperle, D., Krienitz, L., Wolf, M. & Hegewald, E. (2005) Phylogeny of the Hydrodictyaceae (Chlorophyceae): inferences from rDNA data, *Journal of Phycology*, 41, 1039 – 1054.
- CEN (2004) *Water quality – guidance standard for the routine analysis of phytoplankton abundance and composition using inverted microscopy (Utermohl technique)*, CEN/TC230/WG2/TG3.
- CEN (2008) *Water quality – phytoplankton biovolume determination by microscopic measurement of cell dimensions*, CEN/TC230/WG2/TG3.
- Geriš, R. (2004) *Pediastrum privum* (Printz) Hegewald in the Czech Republic, *Czech Phycology, Olomouc*, 4, 63 – 66.
- Hegewald, E. & Schnepf, E. (1979) *Pediastrum privum* (Printz) Hegewald comb. nova, *Algological Studies*, 22, 24 – 28.
- Hegewald E. & Jeon S.L. (2000) The coccobial morphology of *Pediastrum privum* (Printz) Hegewald, *Algological Studies*, 98, 43 – 48.
- Hindák F. & Hindáková A. (2008) Morphology and taxonomy of some rare chlorococcalean algae (Chlorophyta), *Biologia*, 63, 781 – 790.
- Jankovská, V. & Komárek, J. (2000) Indicative value of *Pediastrum* and other coccal green algae in palaeoecology, *Folia Geobotanica*, 35, 59 – 82.
- Komárek, J. & Jankovská, V. (2001) Review of the green algal genus *Pediastrum*; Implication for pollen-analytical research, *Bibliotheca Phycologica*, Band 108, 127., Gebrüder Borntraeger Verlagsbuchhandlung, Berlin, Stuttgart.
- Kowalska, J. & Wołowski, K. (2010) *Pediastrum privum* (Printz) Hegewald (Chlorophyceae) in Lake Małe Zmarle, northern Poland, *Oceanological and Hydrobiological Studies*, 39, 137 – 143.
- Negro, A.I., De Hoyos, C. & Vega, J.C. (2000) Phytoplankton structure and dynamics in Lake Sanabria and Valparaíso reservoir (NW Spain), *Hydrobiologia*, 424, 25 – 37.
- Pełechaty M., Pełechata A., Pukacz, A. (2007) Charophyte flora and vegetation against the background of the trophy state of lakes of Lubuskie Lakeland (mid-Western Poland), Bogucki Wydawnictwo Naukowe, Poznań, 137.
- Prescott G.W. (1962) *Algae of the Western Great Lake Area*, W.C. Brown Company Publishers, Dubuque, Iowa.
- Printz, H. (1914) Kristianiatraktens Protococcoideer, Skrifter Utgit av Videnskapselskapet i Kristiania, *Matematisk-Naturvidenskabelig Klasse*, 1913, 1–121.
- Rojó, C., Segura, M., Rodrigo, M.A. & Salazar, G. (2008) Factors controlling the colonial structure of *Pediastrum tetras* (Chlorophyceae), *Hydrobiologia*, 617, 143 – 155.
- Smith, G.M. (1920) Phytoplankton of the Inland Lakes of Wisconsin, Part I, *Bulletin of the Wisconsin Geological and Natural History Survey*, 57, 1 – 243.
- Tsarenko, P.M. & John, D.M. (2011) Phylum Chlorophyta (Green Algae) Order Sphaeropleales p. 461-465 In; John, D.M., Whitton, B.A. & Brook, A.J., *The Freshwater Algal Flora of the British Isles*, 2<sup>nd</sup> Edition, Cambridge University Press.

---

## First records of the pygmy sperm whale, *Kogia breviceps*, in Scotland

Andrew C. Kitchener<sup>1</sup>, Jerry S. Herman<sup>1\*</sup>, Robert J. Reid<sup>2</sup> and Neil Anderson<sup>3</sup>

<sup>1</sup> Department of Natural Sciences, National Museums Scotland, Chambers Street, Edinburgh EH1 1JF

<sup>2</sup> SAC Consulting: Veterinary Services, Drummondhill Stratherrick Road, Inverness, IV2 4JZ

<sup>3</sup> Houser, Tingwall, Shetland ZE2 9SF

E-mail: a.kitchener@nms.ac.uk

\*E-mail: j.herman@nms.ac.uk

---

The pygmy sperm whale, *Kogia breviceps*, is a poorly known cetacean species, which has been recorded rarely in the British Isles (Leaper and Evans, 2008). It is an oceanic species that inhabits tropical to warmer temperate waters worldwide (Caldwell and Caldwell, 1989). In the North Atlantic it strands reasonably commonly on the coast of the southeast USA (125 strandings between Puerto Rico and Maine 1999-2003 (Waring *et al.*, 2005) as far north as Canada, and in the eastern Atlantic it has been recorded from the Bay of Biscay, stranding from Portugal to the western coast of France with fewer records from the Netherlands and the British Isles (Evans, 1991; Santos *et al.*, 2006). Here we record the first strandings of pygmy sperm whales in Scotland.

Pygmy sperm whales are usually found in small groups of up to six individuals, but more often they are seen alone or in pairs; strandings are most often large males, or mothers and their calves of varying ages, or single females that have recently given birth (Caldwell and Caldwell, 1989; McAlpine, 2002). They feed mostly on